

UDK 616.441-02:616.72-002.77

The principles of timely diagnosis of thyroid pathology in patients with rheumatoid arthritis and autoimmune thyroiditis

Isamitdinova Nilufar Shakhobovna

Bukhara Regional Multidisciplinary Medical Center, Republic of Uzbekistan,
Bukhara city

Email address : nelya--89@list.ru

Resume : Rheumatoid arthritis is a systemic autoimmune rheumatic disease characterized by inflammation of the synovial membranes of the joints, progressive destruction of cartilage and bone tissue, and a wide spectrum of non-joint organ damage[8]. The social importance of RA is the early development of high-level disability. AIT occurs in 2% of the working-age population and is more common in women, leading the cluster of other co-occurring autoimmune diseases (rheumatoid arthritis, type 1 diabetes, vitiligo, etc.). The problem of diagnosis of RA and AIT in the world has not been fully resolved until now, and a number of scientific researches are being carried out in order to practically study the specific aspects of accurate diagnosis in its early period. In this regard, it is necessary to evaluate the impact of both diseases on the main disease outcome based on the identification of risk factors, clarify the changes of RA pathology according to the age of the patients, the duration of the disease activity, clinical course and the effect of drug treatment, as a result of the development of these two diseases, human functional limitations and low quality of life to the fact that it causes severe disability, it is especially important to develop a proper rehabilitation strategy for people suffering from a number of co-morbidities.

Key words: thyroid pathology, rheumatoid arthritis, autoimmune thyroiditis, thyroid gland, hypothyroidism

Introduction. The principles of treating patients with rheumatoid arthritis and autoimmune thyroiditis are one of the complex problems of modern rheumatology. Its relevance is the progression of two autoimmune diseases (RA and AIT), the severity of damage to the musculoskeletal system, the high number of injuries among the working population, early decline in functional ability, loss of professional and social skills, difficulty in physical and psychological adaptation in patients with loss of motor function. It is defined as a general medical and social problem that causes significant economic loss, ie disability.

Prolongation and acceleration of the duration of the main disease leads to the development of thyroid pathology, the addition of new joints and extra-articular systems to the pathological process, the formation of irreversible, permanent changes in the joints (destruction, half-extinction, contracture). All this leads to severe functional deficiency, disruption of all aspects of life and a decrease in the quality of life of patients [3,5,7,12,25,26,27,28,29]. The high medical and social importance of RA and AIT among autoimmune diseases is due to its widespread and early disability and reduced life expectancy of patients [1,4,8,9,10,11,17,21,22].

The diagnosis of RA is still based on a set of clinical and laboratory indicators in the form of constantly changing classification criteria. The visceral manifestations of RA stand out because they determine not only the severity and outcome of the disease, but also its outcome. RA shortens patients' life expectancy by an average of 10 years, causes of death are generally similar to those in the general population, but early cardiovascular and renal disease, pulmonary and gastrointestinal infections are high among the causes of death [2,3,6,18,20,23,24,25,26,27,28,29,30]. Currently, RA is a chronic systemic inflammatory disease that affects not only joint and peri-articular tissues, but also central and autonomic nervous system, endocrine systems and their interrelationships and further complications of the course of the main disease [31,32,33,34,35]. Clinicians are familiar with pathologies such as goitre, hypothyroidism, adrenal gland dysfunction, which are common pathologies of endocrine glands in patients with RA. Thyroid hormones increase the activity of metabolic processes, stimulate lipogenesis, enhance glucose uptake by muscle and adipose tissue, and activate glucogenesis and glyconeogenesis [1]. Thyroid hormones (triiodothyronine-T3, thyroxine-T4) enhance the resorption and synthesis of bone tissue, the production of connective tissue protein glycans and glycosomes. An increase in their amount increases the number of growing and active osteoclasts, accelerates the metabolism of bone tissue, and stimulates osteoblastic function, and this process increases bone tissue-forming markers in the blood [1].

In the absence of thyroid hormones, the amount of adenyl cyclase in the synovial membrane increases, fibroblasts increase the production of hyaluronic acid, which in turn leads to the accumulation of synovial fluid in the joints and causes synovitis [4]. In clinical practice, tests for TTG receptors and antiperoxidase antibodies are carried out in the tissue of thyroid gland [2].

The high frequency of thyroid pathology (42.5%) in patients with RA prompted us to study in detail the processes of the formation of antibodies to thyroid hormones, which depend on the activity and form of RA.

Purpose: In patients with rheumatoid arthritis and autoimmune thyroiditis timely diagnosis of thyroid pathology and development of measures to prevent complications.

Research material and methods: The study was conducted on clinical, instrumental, laboratory analyzes of all patients with rheumatoid arthritis (RA) treated inpatient at BRMMC (Bukhara Regional Multidisciplinary Medical Center) rheumatology department during 2018 and 2020, and among them 82 RA patients (42 RA and 40 RA+AIT) patients were prospectively and 20 healthy controls were selected. The diagnosis of RA was made based on the diagnostic criteria based on the American Society of Rheumatology 2010 classification.

In the first empirical stage, only patients with rheumatoid arthritis, autoimmune thyroiditis and a healthy control group matched for age and gender were selected. A total of 82 patients were included in the study, including 42 patients with rheumatoid arthritis alone, 40 patients with rheumatoid arthritis and autoimmune

thyroiditis, and 20 healthy controls. The course of patients' main disease, concomitant disease, as well as drug anamnesis and pharmacotherapy were evaluated. All patients underwent complex hematological, biochemical, thyroid hormones and anti-antibodies, thyroid ultrasound examination, and densitometry examinations that evaluated all manifestations of RA and AIT accepted in comprehensive rheumatological clinical practice. Hemoglobin, general urinalysis, serum creatinine, calcium, phosphorus, urea, albumin, ALT, AST, T3, T4, TSG (thyroid stimulating hormone), antiTPO, antiTG were performed laboratory tests.

Table-1
Description of the rheumatoid arthritis and control group selected for the study

Indicator	RA (n-82)	Control group (n-20)	R
Gender (female %)	76(92.6%)	16(82%)	0.62
Age	50.58±1.6	48.45 ± 2.1	0.6
Villagers	55(67%)	13(65%)	0.7
Hypertension	37(46%)	9 (43%)	0.04
Duration of illness, year	8.82±1.12	-	-
Type 2 diabetes	6(7%)	1(5%)	0.05
Type of therapy:	82 (100%)	-	-
Methotrexate	76 (92.6%)	-	-
Glucocorticoids	28 (35.2%)	-	-
Leflunomide	13 (15.7%)	-	-
NSAIDs	80 (97.2%)	-	-
GEBT	3 (3.7%)	-	-
Combined therapy	45 (54.9%)	-	-

During the study, 76 (92.6 %) and 16 (82 %) women, 55 (67 %) and 13 (65 %) rural residents, and 37 (46 %) hypertensive patients in the 82 RA patients and 20 control groups, respectively. and 9 (43%), type 2 diabetes was 6 (7%) and 1 (5%). 76 (92.6 %) of patients with RA received methotrexate, 28 (35.2 %) glucocorticosteroids, 13 (15.7 %) leflunomide, 80 (97.2 %) NSAIDS, 3 (3 %) GEBT (genetic engineering basic therapy), 45 (54, 9 %) received combination therapy. Thus, during the study, the majority of patients with rheumatoid arthritis were taking NSAIDS and methotrexate drugs continuously.

At the next stage of the study, 82 patients with rheumatoid arthritis were divided into two groups - 42 with only rheumatoid arthritis and 40 with rheumatoid arthritis combined with autoimmune thyroiditis, based on the results of questionnaires, clinical and laboratory-instrumental examinations.

Table 2

Description of RA and RA+AIT patients

Indicator	RA (n=42)	RA+AIT (n=40)	n-82(100%)
Women, n (%)	38(90.4%)	38(95.0%)	76(92.6%)
Men, n (%)	4(9.5%)	2(5.0%)	6(7.4%)
Average age , years	50.73±1.30 let	53.62±1.48	51.6±1.34
Body mass index , kg/m2	25.56±0.24	25.34±0.28	25.46±0.25
Duration of rheumatoid arthritis , years	8.77±1.12	9.0±1.10	8.88±1.12
RF (+), n (%)	35 (83%)	32 (80%)	67(81.7%)
ACCP (+), n (%)	11 (26.6%)	22 (55%)	33(40.2%)
DAS-28, scores	5.4±0.15	5.68±0.15	5.55±
Process activity :			
Level I (low)	10(23.8%)	5 (12.5 %)	15 (18.3 %)
Level II (medium)	22(52.3%)	20(50%)	42(51.2%)
Level III (higher)	10(23.8%)	15 (37.5 %)	25 (30.5 %)
Functional class , n (%)			
I	2 (4.8%)	4 (10.7%)	6(7.4%)
II	8 (19.0%)	7 (17.5%)	15(18.3%)
III	20 (47.6%)	28 (70.0%)	58(70.7%)
V	2 (4.8%)	1 (2.5%)	3(3.6%)
X-ray stages (according to Steinbroker), n (%)			
I	4(12%)	3 (7.5%)	7 (8.5 %)
II	24 (57.1 %)	16 (45%)	40 (48.7 %)
III	12 (26.2 %)	17 (42.5%)	29 (35.3 %)
IV	2 (4.7%)	4 (5%)	6 (7.5 %)
The presence of extra-articular (systemic) symptoms, n (%)	6 (14.3%)	7 (17.5%)	13(17.0%)
Patients receiving basic therapy, n (%)	42(100%)	40(100%)	82(100%)

Based on the analysis of the table, it was found that there was almost no difference between rheumatoid arthritis (82) and control group (20) by age, gender, comorbidities and region of residence. 82 patients, 76 women (92.6%) and 6 men (7.4%) treated with RA in the rheumatology department of the Bukhara regional multidisciplinary medical center were examined. Among them, patients with rheumatoid arthritis, 55 women (72.4%) and 4 men (25%) belonged to the socially active population, and their age group exceeded 55 to 60 years, respectively. Thus, it was found that the majority of patients with RA were of working age, 50 (61%). The average age of the patients was 50.58 ± 1.34 years. The average duration of the disease was 8.88 ± 1.12 years.

In addition, during the study, all patients filled out a questionnaire identifying thyroiditis and hypothyroidism. Application of questionnaires plays an important role for early diagnosis of the disease and prevention of complications.

Table 3**Questionnaire for the diagnosis of hypothyroidism**

Symbol	Yes, there is	It lasts a long time	No
Unexplained disorders of gastrointestinal functions (mainly constipation)	+40	+6	-5
Weight gain	+25	+5	-3
Abnormal drying of the skin (without cosmetics, climate change)	+30	+6	-15
Fatigue, weakness	+4	+3	-13
Daytime sleepiness	+26	+5	-6
Increased hair loss	+20	+5	-3
Frequent occurrence of swelling (in the previous drinking mode)	+20	+4	-19
Decreased voice timbre, hoarseness (in non-smokers, respiratory system diseases are excluded).	+39	+4	-6

Any total score greater than 7 points from the questionnaire will serve as a basis for referral to an endocrinologist.

Table 4**Symptoms identified on the basis of questionnaires**

Symptoms _	RA+AIT		RA	
	N (40)	%	N (42)	%
Increase in swelling	30	75	14	33
Drowsiness	24	60	11	26
Delayed reaction	30	75	20	54
Increased fatigue	28	70	13	30
Memory impairment	22	55	19	45
Apathy and low mood	15	38	12	29
Bradycardia	12	30	8	20
Weight gain	35	30	7	18
Hair loss	28	70	13	32
Skin dryness	30	75	13	30
Constipation	30	75	6	15

The analysis of the questionnaires showed that before the examination of the

thyroid status, it is possible to identify the symptoms of thyroid gland pathology in patients early and to make a plan for further targeted examination of patients.

As a result of the study, rheumatoid arthritis patients with autoimmune thyroiditis had dry skin in 30 (75%), constipation, slowed reaction, increased fatigue in 28 (70%), increased body weight, drowsiness in 24 (60%), memory impairment in 22 (55%) It was found that only those with rheumatoid arthritis had a relatively high percentage of the mentioned symptoms.

RA was diagnosed according to the American College of Rheumatology (ACR) 1987 diagnostic criteria and the 2010 ACR/EULAR classification [33]. Pain intensity was assessed using a 100 mm visual analog scale (VAS). The DAS-28 index was used to evaluate RA activity according to modern EULAR recommendations. RA is in disease remission when the DAS-28 index is <2.6 , moderately active when the DAS-28 is <3.2 ($3.2 < \text{DAS-28} < 5.1$), and disease active when the DAS-28 index is >5.1 was rated as high. The functional class was evaluated based on the following criteria: I - self-service, professional and non-professional activity ability is maintained; II - the ability to self-service, professional activity is preserved; there is limited non-professional activity; III - self-service ability is preserved, non-professional and professional activities are limited; IV - self-service, professional and non-professional activities are limited.

Table-5
Clinical manifestations of symptoms of all examined patients

Symptoms	RA+AIT n=40		RA n=42	
	n	%	n	%
Swollen joints	40	100	31	75
Painful joints	40	100	35	84
Deformation of joints	24	60	16	40
Morning effort	40	100	42	100
Weight gain	28	70	13	30
Dry skin	22	55	19	45
Symptoms of hypothyroidism	9	22	0	0
Bradycardia	12	30	8	20

Table 6
Social characteristics of the examined groups

Indicator	Gender (E / A , %)	Age	Functional class	Disability , n (%)
-----------	------------------------	-----	---------------------	--------------------

RA (n=42)	4/39 (8%-92%)	50.83±1.60 years old	I: 1 (3%) II: 11 (28%) III: 29 (66%) IV: 1 (3%)	13 (16%)
RA+AIT (n=40)	2/37 (9%-91%)	52.92±1.60 years	I: 0 (0 %) II: 4 (11%) III: 34 (85%) IV: 2 (4%)	36 (90%)

According to the table, patients in both groups participating in the study did not differ from each other in terms of gender, age and functional classes of the disease. At the time of inclusion in the study, the number of disabled patients associated with rheumatoid arthritis and autoimmune thyroiditis was several times higher than those with only rheumatoid arthritis ($r > 0.05$). If the II-radiological stage predominates only in patients with RA, destructive processes and ankylosis of joints, characteristic of reliable stages III-IV, were observed in patients with RA combined with AIT.

Only in patients with RA, a significant decrease in functional activity was rarely noted, and most patients continued their professional activities and led an active lifestyle, while patients with RA associated with autoimmune thyroiditis became unable to work in the early stages of the disease.

Table 7 shows data on the disability of both groups of patients

Table 7

The structure of disability

Indicator	disability , n (%)	Group I disability , n (%)	Group II disability , n (%)	Group III disability , n (%)
RA (n=42)	13(31 %)	1 (2 %)	12 (29%)	0 (0%)
RA+AIT (n=40)	14(35 %)	2(5%)	12 (30 %)	0 (0%)

In both groups of patients, mainly II disabilities were recorded, but in patients with rheumatoid arthritis and autoimmune thyroiditis, the number of I disabilities was 2 times higher than in patients with only rheumatoid arthritis.

The level of activity of the pathological process was determined using diagnostic criteria, working classification and DAS 28 (Disease Activity Score) index. 9 (11.1%) patients had a score of 3.2 or less on the DAS 28 index, corresponding to the I activity level of RA, and 44 patients with a DAS 28 index of 3.2 to 5.1 (activity level II) 53.7%) patients and DAS 28 index higher than 5.1 (level of activity III) – 29 (35.2%) patients. According to the DAS-28 index,

similar activity levels were found in the patients, and in both groups this indicator showed higher than average RA activity level results. Seropositive RA was detected in 67 patients, and seronegative - in 15 patients. In RA with AIT, the seropositive type was reliably detected more often (81.7%, compared to 18.3% in the control group), and in most cases it was severe.

Summary

The principles of treatment of patients with rheumatoid arthritis and autoimmune thyroiditis are one of the complex problems of modern rheumatology. Its relevance is the progression of two autoimmune diseases (RA and AIT), the severity of damage to the musculoskeletal system, the high number of injuries among the working population, early decline in functional ability, loss of professional and social skills, difficulty in physical and psychological adaptation in patients with loss of motor function. It is defined as a general medical and social problem that causes significant economic loss, i.e. disability.

Prolongation and acceleration of the duration of the main disease leads to the development of thyroid pathology, the addition of new joints and extra-articular systems to the pathological process, the formation of irreversible, permanent changes in the joints (destruction, half-extinction, contracture). All this leads to severe functional deficiency, disruption of all aspects of life and a decrease in the quality of life of patients. Given the above reasons and the high prevalence of thyroid gland damage in RA patients, it is reasonable to introduce active detection of signs of subclinical thyroid damage. The use of questionnaires in determining thyroid pathology in patients with RA and AIT will provide practical help to general practitioners, endocrinologists and rheumatologists in making a prognosis of the course of the underlying disease.

List of references

1. Авдеева А.С., Новиков А.А., Александрова Е.Н. и др. Связь уровней цитокинов с активностью заболевания, уровнем аутоантител и деструктивными изменениями суставов при раннем ревматоидном артрите. Научно-практическая ревматология. 2015;53(4):385-90.
2. Агаки С.А., Граппа Л.Г. Взаимосвязь между функцией щитовидной железы и течением ревматоидного артрита // Научно практическая ревматология. – 2001. – № 3. – С. 4
3. науки и образования, 2021. №11 (114). Часть 1. с.13-17
4. Елисеева Л.В. Течение ревматоидного артрита, сочетающегося с патологией щитовидной железы : автореф. дис. ... канд. мед. наук. – Томск, 2002. – 21 с
5. Зборовский А.Б., Фофанова Н.А. Ревматические болезни и ревматическая служба в Южном Федеральном округе: состояние и перспективы. Научно-практическая ревматология. 2007; 3: 4-9.

6. Зиннетулина Н.Х. Особенности тиреоидного статуса у больных ревматоидным артритом : автореф. дис. ... канд. мед. наук. – Чебоксары, 2003. – 23 с.
7. Ajjan R.A., Watson P.F., McIntosh R.S., Weetman A.P. Intrathyroidal cytokine gene expression in Hashimoto's thyroiditis. Clin Exp Immunol. 1996 Sep;105(3):523-8. doi: 10.1046/j.1365- 2249.1996.d01-784.x ;
8. Ajjan R.A., Weetman A.P. Cytokines in thyroid autoimmunity. Autoimmunity. 2003 SepNov;36(67):3519
9. Al-Awadhi A.M, Olusi S., Hasan EA, Abdullah A. Frequency of abnormal thyroid function tests in Kuwaiti Arabs with autoimmune diseases. Med Princ Pract. 2008;17(1):61-5.
10. Al Girgawy Sh. A.F., AlShabrawy D.A. Thyroid disorders and autoantibodies in systemic lupus erythematosus // Egypt. Rheumatol. Rehab. – 2007. – Vol. 34, No. 3. – P. 497–508
11. Хамдамов И.Б. Клиническая оценка эффективности традиционного подхода лечения грыж передней брюшной стенки у женщин фертильного возраста // Вестник врача. –Самарканд 2022. № 2.2 (104).-С.65-70.
12. Khamdamov I.B., Khamdamov A.B. Differentiated approach to the choice of hernioplasty method in women of fertile age (Clinical and experimental study) // Тиббиётда янги кун. – Бухоро, 2021.-№ 6 (38/1).-С. 112-114.
13. Хамдамов И.Б., Хамдамов А.Б. Фертил ёшдаги аёлларда эндовидеохирургик герниопластика // Тиббиётда янги кун. Бухоро, 2021.-№6 (38/1) -С. 25-27.
14. Хамдамов И.Б. Experimental determination of the extensibility of the anterior abdominal wall tissues at different times of pregnancy using various approaches to hernioplasty// Academica: An International Multidisciplinary Research Journal Vol. 12, Issue 04, April 2022 SJIF 2022 = 8.252 P.193-201
15. Хамдамов И.Б. Совершенствование тактических подходов в лечении грыж передней брюшной стенки у женщин фертильного возраста // Тиббиётда янги кун. Бухоро, 2022.-№10(48)- С. 338-342.
16. Хамдамов И.Б. Морфофункциональные особенности брюшного пресса у женщин репродуктивного возраста // Тиббиётда янги кун. Бухоро, 2022.-№3(41)- С. 223-227.
17. Khamdamova M.T. Ultrasound features of three-dimensional echography in assessing the condition of the endometrium and uterine cavity in women of the first period of middle age using intrauterine contraceptives

- // Biology va tibbiyot muammolari. - Samarkand, 2020. - No. 2 (118). - P.127-131.
18. Khamdamova M. T. Ultrasound assessment of changes in the endometrium of the uterus in women of the first and second period of middle age when using intrauterine and oral contraceptives // Биомедицина ва амалиёт журнали. – Ташкент, 2020. - №2. - 8 часть. - С.79-85.
 19. Khamdamova M. T. Anthropometric characteristics of the physical status of women in the first and second period of middle age // A new day in medicine. Tashkent, 2020. - № 1 (29). - С.98-100.
 20. Khamdamova M.T. Age-related and individual variability of the shape and size of the uterus according to morphological and ultrasound studies // News of dermatovenereology and reproductive health. - Tashkent, 2020. - No. 1-2 (88-80). - P.49-52.
 21. Khamdamova M. T. Anthropometric characteristics of the physical status of women in the first and second period of middle age // Тиббиётда янги кун. Ташкент, 2020. - № 1 (29). - С.98-100.
 22. Хамдамова М.Т. Возрастная и индивидуальная изменчивость формы и размеров матки по данным морфологического и ультразвукового исследований // Новости дерматовенерологии и репродуктивного здоровья. - Ташкент, 2020. - № 1-2 (88-80). - С.49-52.
 23. Хамдамова М.Т. Ультразвуковые особенности трехмерной эхографии в оценке состояния эндометрия и полости матки у женщин первого периода среднего возраста применяющие внутриматочные контрацептивные средства // Биология ва тиббиёт муаммолари. - Самарканд, 2020. - №2 (118). - С.127-131.
 24. Khamdamova M. T. Ultrasound assessment of changes in the endometrium of the uterus in women of the first and second period of middle age when using intrauterine and oral contraceptives // Биомедицина ва амалиёт журнали. – Ташкент, 2020. - №2. - 8 часть. - С.79-85.
 25. Хамдамова М.Т. Особенности ультразвуковых параметров матки у женщин первого и второго периода среднего возраста применяющие инъекционные контрацептивные средства // Тиббиётда янги кун. - Ташкент, 2020. - № 2/1 (29/1). - С.154-156.
 26. Хамдамова М.Т. Особенности ультразвукового изображения матки и яичников у женщин второго периода среднего возраста применяющие комбинированные оральные контрацептивные средства // Тиббиётда янги кун. - Ташкент, 2020. - № 2 (30). - С. 258-261.

27. Хамдамова М.Т. Индивидуальная изменчивость матки и яичников у женщин применяющие и не использующие различные виды контрацептивные средства // Тиббиётда янги кун. - Ташкент, 2020. - № 3 (31). - С. 519-526.
28. Khamdamova M. T. Echographic features variability in the size and shape of the uterus and ovaries in women of the second period of adulthood using various contraceptives // Asian Journal of Multidimensional Research - 2020. – N9 (5). - P.259-263.
29. Khamdamova M. T. Somatometric characteristics of women of the first and second period of adulthood using different contraceptives with different body types // The american journal of medical sciences and pharmaceutical research - 2020. – N8 (2). - P.69-76.
30. Хамдамова М.Т., Жалолдинова М.М.,Хамдамов И.Б. Состояние оксида азота в сыворотке крови у больных кожным лейшманиозом // Тиббиётда янги кун. - Бухоро, 2023. - № 5 (55). - С. 638-643.
31. Хамдамова М.Т., Жалолдинова М.М.,Хамдамов И.Б. Значение церулоплазмина и меди в сыворотки крови у женщин носящих медьсодержащих внутриматочной спирали // Тиббиётда янги кун. - Бухоро, 2023. - № 6 (56). - С. 2-7.
32. Khamdamova M. T. Bleeding when wearing intrauterine contraceptives and their relationship with the nitric oxide system // American journal of pediatric medicine and health sciences Volume 01, Issue 07, 2023 ISSN (E): 2993-2149. P.58-62
33. Khamdamova M. T. The state of local immunity in background diseases of the cervix // Eurasian journal of medical and natural sciences Innovative Academy Research Support Center. Volume 3 Issue 1, January 2023 ISSN 2181-287X P.171-175.
34. Хамдамова М.Т., Хасанова М.Т. Различные механизмы патогенез гиперплазии эндометрия у женщин постменопаузального периода (обзор литературы) // Тиббиётда янги кун. - Бухоро, 2023. - № 8 (58). - С. 103-107.
35. Barton A, Worthington J. Genetic susceptibility to rheumatoid arthritis: an emerging picture. Arthritis Rheum. 2009 Oct 15;61(10):1441-6. doi: 10.1002/art.24672
36. Mirabdullayev I. M. et al. New records of *Microcyclops afghanicus* Lindberg, 1948 from Kazakhstan, Central Asia (Copepoda, Cyclopoida) //Crustaceana. – 1997. – Т. 70. – №. 8. – С. 849-854.
37. Рахматуллаева Г. К. и др. COVID-19 АССОЦИИРОВАННЫЙ ТРОМБОЗ КАВЕРНОЗНОГО СИНУСА.(КЛИНИЧЕСКИЙ СЛУЧАЙ) //ЖУРНАЛ НЕВРОЛОГИИ И

НЕЙРОХИРУРГИЧЕСКИХ ИССЛЕДОВАНИЙ. – 2021. – №. SPECIAL 1.

38. Khalimova K. M. et al. Clinical neurological and diagnostic aspects of headaches with congenital precerebral angiodyplasia //Toshkent tibbiyot akademiyasi axborotnomasi. – 2019. – №. 2. – С. 132-134.
39. Salokhiddinov M., Rahmatullaeva G. The role of pathological deformations of the main arteries of the brain In the development< bold> of stroke</bold> //CEREBROVASCULAR DISEASES. – ALLSCHWILERSTRASSE 10, CH-4009 BASEL, SWITZERLAND : KARGER, 2018. – Т. 45. – С. 473-473.
40. Муминова З. А., Нишанов Д. А. Морфологическая характеристика пуповины при беременности, осложненной острой респираторной инфекцией //Журнал теоретической и клинической медицины. – 2016. – №. 4. – С. 104-107.
41. Муминова З. А., Саиджалилова Д. Д., Нишанов Д. А. Морфологическая характеристика плацент женщин, перенесших острые респираторные инфекции //Журнал теоретической и клинической медицины. – 2016. – №. 3. – С. 148-150.
42. Ayupova F. et al. 137. The influence of acute respiratory infection in pregnant women on the development of preeclampsia //Pregnancy Hypertension. – 2018. – Т. 13. – С. S83.

