

SEVERE ANEMIA: ETIOLOGY, PATHOPHYSIOLOGY AND CLINICAL SIGNIFICANCE

Nodira I. Zakirova, Professor

Fotima I. Zakirova, PhD, Associate Professor

Otabek A. Umarov, PhD Student

Department of Obstetrics and Gynecology No. 1, Samarkand State Medical University, Samarkand, Uzbekistan

Abstract

Severe anemia is a complex hematological condition characterized by markedly reduced hemoglobin concentration or red blood cell count, leading to decreased oxygen delivery to tissues. As a major global health concern, severe anemia significantly increases the risk of morbidity and mortality, particularly among vulnerable populations such as pregnant women, children, and individuals with chronic diseases. This article analyzes the etiology, pathophysiological mechanisms, clinical manifestations, diagnostic approaches, treatment strategies, and public health implications of severe anemia. The review highlights the multifactorial nature of the disease and underscores the importance of integrated prevention, early diagnosis, and appropriate therapeutic interventions to reduce its global burden.

Keywords: severe anemia, hemoglobin deficiency, erythropoiesis, hypoxia, micronutrient deficiency, public health

Introduction

Anemia represents one of the most widespread hematological disorders globally, with the World Health Organization (WHO) estimating that approximately one-third of the world's population is affected to varying degrees. Severe anemia, as the most advanced stage of the condition, poses significant clinical and public health challenges. Defined by substantially reduced hemoglobin levels that compromise oxygen delivery and cellular metabolism, severe anemia is associated with systemic physiological dysfunction, reduced productivity,

increased maternal and perinatal complications, and elevated mortality in both children and adults.

The global prevalence of severe anemia is influenced by socio-economic, environmental, nutritional, and pathological factors. Sub-Saharan Africa and South Asia record notably high burdens due to nutritional deficiencies, parasitic infections, hemoglobinopathies, and limited access to healthcare. In industrialized regions, severe anemia is more frequently related to chronic diseases, inherited blood disorders, and malignancies. These regional differences underscore the multi-dimensional nature of severe anemia and the necessity of contextualized health policies and targeted interventions.

The increasing attention to severe anemia in recent decades is linked to its direct impact on developmental outcomes, cognitive performance, maternal health, and workforce productivity. As such, comprehensive scientific understanding of severe anemia is essential for clinicians, researchers, policymakers, and global health programs.

Etiology of Severe Anemia

The etiology of severe anemia is multi-factorial and can be classified into nutritional, genetic, pathological, and environmental categories. Understanding these determinants is crucial for accurate diagnosis and targeted treatment.

Nutritional etiologies remain the leading global cause of anemia. Deficiencies in key micronutrients—particularly iron, vitamin B12, and folate—alter erythropoiesis and hemoglobin synthesis. Iron deficiency impairs heme formation, leading to microcytic and hypochromic anemia. Vitamin B12 and folate deficiencies disrupt DNA synthesis, producing megaloblastic anemia characterized by large, fragile erythrocytes.

Foods lacking in animal protein, leafy greens, and fortified cereals contribute to nutritional anemia, with pregnant women, adolescents, and young children particularly vulnerable due to increased physiological demands.

Chronic diseases such as renal failure, autoimmune disorders, cancer, HIV/AIDS, and inflammatory bowel disease contribute to anemia of chronic

disease (ACD). Mechanisms include suppressed erythropoietin production, shortened red blood cell lifespan, and impaired iron mobilization due to inflammatory cytokines. Patients with end-stage renal disease frequently develop severe anemia without therapeutic intervention.

Genetic disorders such as sickle cell disease and thalassemia directly alter hemoglobin structure and red cell stability. Sickle cell anemia induces hemolysis due to polymerization of abnormal hemoglobin S under hypoxic conditions. Alpha and beta thalassemias reduce globin chain synthesis, producing severe microcytic anemia and requiring consistent medical support.

Acute or chronic hemorrhage significantly lowers circulating hemoglobin. Traumatic injury, gastrointestinal bleeding, heavy menstrual bleeding (menorrhagia), or post-surgical complications may induce severe anemia in the absence of timely intervention. Chronic occult bleeding is common in peptic ulcer disease, colorectal malignancies, and parasitic infections.

Bone marrow suppression impairs hematopoiesis and may result from aplastic anemia, myelodysplastic syndromes, malignancies, chemotherapy, radiation exposure, and viral infections (e.g., Parvovirus B19). These conditions reduce red blood cell production and contribute to severe anemia despite adequate iron stores.

The pathophysiological mechanisms of severe anemia are primarily linked to tissue hypoxia resulting from reduced hemoglobin concentration and impaired oxygen transport capacity. Compensatory physiological responses aim to preserve oxygen delivery and cellular function.

Hemoglobin molecules bind oxygen in the lungs and transport it to tissues. Severe hemoglobin reduction disrupts this function, causing systemic oxygen deprivation. Brain and cardiac tissues are particularly sensitive to hypoxia due to high metabolic demands.

To compensate for decreased oxygen-carrying capacity, the cardiovascular system increases cardiac output through tachycardia and stroke volume

augmentation. Prolonged compensation places strain on the heart and may contribute to heart failure, particularly in elderly or chronically ill patients.

Reduced oxygen availability forces cells to shift from aerobic to anaerobic metabolism, increasing lactate production and resulting in metabolic acidosis. Chronic hypoxia disrupts mitochondrial function and contributes to multi-organ impairment.

Cognitive dysfunction, dizziness, and attention deficits result from reduced cerebral perfusion. In children, prolonged severe anemia is associated with delayed cognitive development and impaired academic performance.

Conclusion

Severe anemia is a major health burden with diverse etiological factors, complex pathophysiological mechanisms, and significant clinical consequences. Early diagnosis, effective treatment, and preventive public health strategies can substantially reduce disease burden. Continuous research is required to improve clinical management, strengthen screening programs, and enhance global health outcomes.

REFERENCES

- 1.Ubbiniyazova A. M. Characteristic of a functional condition of an organism of workers of the Tashkent tractor plant //Medical magazine of Uzbekistan. – 2012. – C. 80-82.
- 2.Ataniyazova R. A. et al. Assessment of knowledge and skills of hand hygiene among representatives of three generations of the family living in different regions of Uzbekistan.
- 3.Madaminova M. A. et al. The dynamics are changes in organoleptic parameters and chemical composition of amudarya river water //European Journal of Molecular and Clinical Medicine. – 2020. – T. 7. – №. 7. – C. 537-545.
- 4.Атаханова Д. О., Уббиниязова А. М., Садирова М. К. ОСНОВНЫЕ ГИГИЕНИЧЕСКИЕ ПРОБЛЕМЫ КЛАССИФИКАЦИИ ЗАГРЯЗНЕНИЯ ПОЧВ В КАРАКАЛПАКСТАНЕ //The Way of Science. – 2014. – С. 77.

5. Iskandarova G. T., Ubbiniyazova A. M. PROBLEMS OF ANTI-
INFECTION PROTECTION IN BEAUTY AND HEALTH INSTITUTIONS

// Медицинский журнал молодых ученых. – 2025. – №. 15 (09). – С. 202-205.