

Impact of Metabolic Syndrome on Bone Regeneration during Fracture**Healing****Akromov Iskandar****Abdurakhmanov Mirsaid Mirazizovich****Abstract**

Metabolic syndrome is a widespread health condition characterized by a combination of obesity, insulin resistance, hyperglycemia, dyslipidemia, and hypertension. These metabolic abnormalities have profound effects on various physiological systems, including the skeletal system. Bone regeneration following fractures is a highly coordinated process involving inflammation, angiogenesis, cellular proliferation, and remodeling. The present study examines the impact of metabolic syndrome on bone regeneration during fracture healing. Experimental observations indicate that metabolic syndrome delays callus formation, impairs osteoblast differentiation, reduces vascularization, and prolongs the remodeling phase. Histological and radiological analyses demonstrate significant differences between healthy and metabolically compromised subjects. Understanding these effects is essential for developing targeted therapeutic strategies to improve fracture outcomes in patients with metabolic disorders.

Keywords: Metabolic syndrome, bone regeneration, fracture healing, osteogenesis, insulin resistance, angiogenesis, osteoblasts, bone remodeling.

Introduction

Bone fractures represent one of the most common musculoskeletal injuries worldwide. Under normal physiological conditions, bone healing proceeds through a sequence of overlapping stages, including inflammation, soft callus formation, hard callus development, and remodeling. This process restores both the structural integrity and mechanical strength of the damaged bone.

The prevalence of metabolic syndrome has increased dramatically over the past decades due to sedentary lifestyles, unhealthy dietary habits, and rising obesity rates. Metabolic syndrome is recognized as a major risk factor for

cardiovascular disease and type 2 diabetes mellitus. However, its influence on skeletal health has attracted growing scientific interest.

Bone tissue is metabolically active and highly sensitive to systemic physiological changes. Conditions associated with metabolic syndrome, such as chronic inflammation, oxidative stress, and impaired glucose metabolism, can disrupt the balance between bone formation and bone resorption. Consequently, fracture healing may become delayed or incomplete.

The purpose of this study is to investigate the effects of metabolic syndrome on bone regeneration during fracture repair and to identify the biological mechanisms responsible for impaired osteogenesis.

Materials and Methods

The study was conducted using an experimental animal model. Laboratory rats were divided into two groups: a control group and a metabolic syndrome group. Metabolic syndrome was induced through prolonged consumption of a high-fat, high-sugar diet combined with reduced physical activity.

Following confirmation of metabolic syndrome through biochemical analyses, standardized femoral fractures were created under sterile conditions and general anesthesia. The fractures were stabilized to ensure proper alignment and comparable healing conditions across all animals.

Animals were monitored throughout the healing period. Tissue samples were collected at different stages of fracture repair for histological examination. Radiographic studies were performed to assess callus formation, mineralization, and fracture union.

Histological evaluation focused on osteoblast activity, vascular development, inflammatory cell infiltration, and trabecular bone formation. Comparative analysis was conducted between the experimental and control groups.

Results and Discussion

The findings demonstrated that metabolic syndrome significantly impaired bone regeneration. During the inflammatory phase, animals with metabolic

syndrome exhibited prolonged inflammatory responses characterized by increased infiltration of inflammatory cells and elevated tissue edema.

Soft callus formation occurred later in the experimental group compared with controls. Histological examination revealed reduced proliferation and differentiation of osteoblast precursor cells. As a result, bone matrix synthesis was significantly decreased.

Angiogenesis was also adversely affected. Newly formed blood vessels were less numerous and poorly organized within the fracture site. Since adequate vascularization is essential for delivering oxygen, nutrients, and growth factors, impaired angiogenesis contributed substantially to delayed healing.

Radiographic assessment showed reduced mineral density and slower progression toward fracture consolidation. While the control group demonstrated advanced callus mineralization by the eighth week, the metabolic syndrome group exhibited incomplete ossification and persistent fracture gaps.

Several biological mechanisms may explain these observations. Chronic low-grade inflammation associated with metabolic syndrome increases levels of cytokines such as tumor necrosis factor- α (TNF- α) and interleukin-6 (IL-6), which inhibit osteoblast activity. Furthermore, insulin resistance compromises cellular energy metabolism and reduces the anabolic effects of insulin on bone tissue.

Oxidative stress also plays a critical role by damaging cellular structures and disrupting signaling pathways involved in osteogenesis. Together, these factors create an unfavorable environment for effective bone regeneration.

The results confirm that metabolic syndrome is not only a metabolic disorder but also a significant risk factor for impaired skeletal repair and delayed fracture healing.

Conclusion

Metabolic syndrome exerts a negative influence on bone regeneration during fracture healing. Delayed callus formation, impaired angiogenesis, reduced

osteoblast activity, and prolonged remodeling were identified as key pathological features. These alterations contribute to slower fracture union and increased risk of complications. Effective management of metabolic syndrome and development of targeted regenerative therapies may improve fracture healing outcomes and enhance the quality of life for affected patients. Future studies should investigate molecular pathways involved in osteogenesis under metabolic conditions and explore innovative treatment strategies aimed at restoring normal bone regeneration.

REFERENCES

- 1.Akhtam R. et al. Biomarkers in liver regeneration //Clinica Chimica Acta. – 2025. – T. 576. – C. 120413.
- 2.Shagzatova B. X., Mirkhaydarova F. S. FEATURES OF DIABETES MELLITUS WITH HIV-INFECTED PATIENTS //EDITOR COORDINATOR. – 2021. – C. 778.
- 3.Akhmedova F. et al. The course of Parkinson's disease in patients with impaired carbohydrate metabolism //MOVEMENT DISORDERS. – 111 RIVER ST, HOBOKEN 07030-5774, NJ USA : WILEY, 2018. – T. 33. – C. S176-S177.
- 4.Shagzatova B. X. et al. Endokrinologiya mutaxassisligi bo'yicha klinik rezidentlarni o'qitishda «case»-usuli (Doctoral dissertation, Ўзбекистон, Тошкент). – 2023.
- 5.Шагазатова Б. Х. и др. Особенности течения сахарного диабета у вич-инфицированных больных. – 2019.