

STUDY OF THE CHEMICAL COMPOSITION OF BITTER PLANTS (*Artemisia absinthium L.*).

Toyiraxon Amirova Sheraliyevna

PhD, the senior lecturer, Department of chemistry, Fergana State University

Nishonova Robiyaxon Muhammadzohir qizi

Graduate student of Ferghana State University

Erkinov Jamshidbek Dilshodbek ugli

student of Fergana State University

Abstract: Phenolcarboxylic acids (not less than 0.5% by chlorogenic acid). Coumarins (1.2%): scopoletin, umbelliferone, emperorin, esculetin, xanthotoxol, coumarin. Flavonoids (not less than 0.5% of rutin): quercetin, kaempferol, isorhamnetin, apigenin. Carbohydrates: polysaccharides, inulin, starch.

Key words: scopoletin, umbelliferone, emperorin, esculetin, xanthotoxol, polysaccharides, inulin, starch.

Botanical and morphological-anatomical description of wormwood. Wormwood (*Artemisia absinthium L.*) is a perennial herbaceous plant from the rhizome of which develops several stems 2 meters or more in size.

The arrangement of the leaves is alternate. Stem leaves: petiolate, rounded triangular, twice or thrice pinnatisected; apical: sessile, entire or tripartite, reticulate venation, entire margin, dense pubescence on both sides of the leaf, grayish green. Stem: ribbed, short-branched, with dense grayish-green pubescence.

Flowers: inflorescence - spherical baskets forming a paniculate inflorescence; involucral leaflets are oblong-oval, outer ones are almost the same length as inner ones, concave along the back, broadly membranous along the margin.

Chemical composition. Wormwood has a strong pleasant spicy smell. Wormwood herb contains an essential oil (0.07-0.2%), in its composition: limonene, terpinolene, fenchon, aromadendren, tayyl alcohol, thujone (alpha and betta), alpha-pinene, beta-pinene, camphor, alpha-copaene, alpha-guayene, beta-caryophyllene, borneol, cadinene, vulgarol, tricyclene, camphene, alpha-terpinene, artemisimaketone, artemisian alcohol, isoborneol. Triterpenoids: alpha-amirin, fernol. Steroids: beta-sitosterol, stigmasterol.

Amino acids: arginine, histidine, asparagine, proline, lysine, alanine, valine, glycine, isoleucine, aspartic acid, methionine. Phenolcarboxylic acids (not less than 0.5% in terms of chlorogenic acid).

Coumarins (1.2%): scopoletin, umbelliferon, emperorin, esculetin, xanthotoxol, coumarin. Flavonoids (not less than 0.5% in terms of rutin): quercetin, kaempferol, isorhamnetin, apigenin. Carbohydrates: polysaccharides, inulin, starch.

LITERATURE

- 1.Блинова К. Ф. и др.Ботанико-фармакогностический словарь: Справ. пособие/ Под ред. К. Ф. Блиновой, Г. П. Яковлева-М.: Высш. шк., 1990. -С.226-227.-ISBN 5-06-000085-0.
- 2.К.Ш.Тожибаев, Н.Ю.Бешко, Х.Ф.Шомуродов, У.Х.Кодиров, О.Т.Тургунов, В.К.Шарипова. Кадастр флоры.Узбекистана Кашкадарьинская область. Тошкент-2019.
- 3.A.A.Ibragimov, T. Sh.Amirova, A.A.Ibrokhimov. Certification and classification of tissues based on their biological properties and chemical composition. *Universum: Chemistry and biology: Sci. Jorn.* 2020. № 10 (76), oktober, p.10-14.
- 4.T.Amirova, A.Ibragimov, O.Nazarov. Coloring Natural Silk with Natural Dyes Obtained from Plants. *Annals of the Romanian Society for Cell Biology*, 2021, 7089–7093. Retrieved from <https://www.annalsofrscb.ro/index.php/journal/article/view/2225>.
- 5.Назаров, О. М., & Амирова, Т. Ш. (2022). ОПРЕДЕЛЕНИЕ СОДЕРЖАНИЯ МАКРО-И МИКРОЭЛЕМЕНТОВ В РАЗЛИЧНЫХ ВИДАХ КОЖИ МЕТОДОМ МАСС-СПЕКТРОМЕТРИИ С ИНДУКТИВНО-СВЯЗАННОЙ ПЛАЗМОЙ. Главный редактор, 18.
- 6.Амирова, Т. Ш. (2022, April). ХИМИЧЕСКАЯ ПОДГОТОВКА ТКАНЕЙ ИЗ НАТУРАЛЬНОГО ШЁЛКА. In Conference Zone (pp. 137-138).